

NATIONAL INSTITUTE FOR HEALTH AND CLINICAL EXCELLENCE AND SOCIAL CARE INSTITUTE FOR EXCELLENCE

SCOPE

1 Guideline title

Conduct disorders and associated antisocial behaviour: recognition, intervention and management of conduct disorders and associated antisocial behaviour in children and young people

Short title

Conduct disorders in children and young people

2 The remit

The Department of Health has asked NICE and the Social Care Institute for Excellence (SCIE): 'To produce a clinical guideline on the recognition, identification and management of conduct disorder (including oppositional defiance disorder) in children and young people.'

3 Clinical need for the guideline

3.1 Epidemiology

- a) Conduct disorders are characterised by repetitive and persistent patterns of antisocial, aggressive or defiant behaviour that consists of significant and persistent violations of age-appropriate social expectations. The current World Health Organization classification of the disorders (ICD-10) identifies two subgroups: conduct disorder and oppositional defiant disorder. Conduct disorder is more common in older children (11 to 12 years and older) and oppositional defiant disorder more common in those aged 10 years or younger. The major distinction between the disorders is the extent and the severity of the antisocial behaviour. Isolated

antisocial or criminal acts are not sufficient to support a diagnosis of conduct disorder or oppositional defiant disorder.

- b) Conduct disorders are the most common mental health disorder in children and young people. The Office of National Statistics surveys of 1999 and 2004 reported that prevalence for conduct disorders with associated impairment was 5% among children and young people. The prevalence without impairment was not much larger, because conduct disorders nearly always have a significant impact on functioning and quality of life. The first survey demonstrated that conduct disorders have a steep social class gradient, with a three- to fourfold increase in the lowest social classes compared with the highest. The second survey found that almost 40% of looked after children, those who have been abused and/or those on child protection/safeguarding registers, between 5 and 17 years old, have conduct disorders.
- c) The prevalence of conduct disorders increases throughout childhood and they are more common in boys than girls. For example, 7% of boys and 3% of girls aged 5 to 10 years have conduct disorders; for children aged 11 to 16 years the number rises to 8% for boys and 5% for girls.
- d) Conduct disorders commonly coexist with other mental health disorders, for example, 46% of boys and 36% of girls have at least one other coexisting mental health disorder. The coexistence of conduct disorders with attention deficit hyperactivity disorder (ADHD) is particularly high and in some groups more than 40% of people with a diagnosis of conduct disorder also have a diagnosis of ADHD. The presence of conduct disorder in childhood is also associated with a significantly increased rate of mental health disorders in adult life, including antisocial personality disorder (up to 50% of children and young people with a conduct disorder may go onto develop antisocial personality disorder). The prevalence of conduct disorders varies between different ethnic groups, being

lower than average in some groups (for example, British Indian Children 2.1%) but higher in other groups (for example, Black British children 8.6%)

- e) A diagnosis of a conduct disorder is strongly associated with poor educational performance (children with conduct disorders are five times more likely to leave school with no qualifications), social isolation, drug and alcohol misuse and increased contact with the criminal justice system. This association continues into adult life with poorer educational and occupational outcomes, involvement with the criminal justice system (as high as 50% in some groups) and a high level of mental health disorder (at some point in their lives 90% of people with antisocial personality disorder will suffer from another mental disorder).

3.2 Current practice

- a) Conduct disorders are the most common reason for referral of young children to child and adolescent mental health services (CAMHS). Children with conduct disorders also comprise a considerable proportion of the work of the health and social care system; for example, 30% of a general practitioner's child consultations are for conduct disorders, 45% of community child health referrals are for behaviour disturbances, and psychiatric disorders are a factor in 28% of paediatric outpatient referrals. In addition, social care services have significant involvement with children and young people with conduct disorders, with more vulnerable or disturbed children often being placed with a foster family or, in a small number of cases, in residential care. The demands on the educational system are also considerable and include the provision of special needs education. The criminal justice system also has significant involvement with older children with conduct disorders.
- b) Multiple agencies may be involved in the care and treatment of children with conduct disorders, which presents a major challenge

for current services in the effective coordination of care across agencies.

- c) Several interventions have been developed for children with conduct disorders and related problems. These have been covered in 'Parent-training/education programmes in the management of children with conduct disorders', NICE technology appraisal guidance 102 (2006) and 'Antisocial personality disorder: treatment, management and prevention', NICE clinical guideline 77 (2009). Other interventions focused on prevention, such as the Family Nurse Partnership, have recently been implemented in the UK and are currently being evaluated. Three themes are common to these interventions: a strong focus on working with parents and families, recognition of the importance of the wider social system in enabling effective interventions, and a focus on preventing or reducing the escalation of existing problems. However, the availability of the majority of these interventions varies across the country. Parenting programmes are the best established; implementation of multisystemic approaches and early intervention programmes is more variable. In addition to the programmes developed specifically for children with conduct disorders, a number of children (and their families) are treated by both specialist CAMHS teams and general community-based services such as Sure Start. Identifying which of the above interventions and agencies are the most appropriate remains challenging especially for non-specialist health, social care and educational services. Further challenges arise when considering the use of prevention and early intervention programmes and identifying which vulnerable groups stand to gain from such interventions. Factors that may be associated with a higher risk of developing conduct disorders include parental factors such as parenting style and parental mental health and other problems (such as antisocial personality disorder and substance misuse); environmental factors such as

poverty and place of residence (for example, foster care); and the presence of other mental disorders such as ADHD.

4 The guideline

The guideline development process is described in detail on the NICE website (see section 6, 'Further information').

This scope defines what the guideline will (and will not) examine, and what the guideline developers will consider. The scope is based on the referral from the Department of Health.

The areas that will be addressed by the guideline are described in the following sections.

4.1 Population

4.1.1 Groups that will be covered

- a) Children and young people (18 years and younger) with a diagnosed or suspected conduct disorder, including looked after children and those in contact with the criminal justice system.
- b) Children and young people identified as being at significant risk of developing conduct disorders.
- c) Consideration will be given to the specific needs of:
 - children and young people with conduct disorders and coexisting conditions (such as ADHD, depression, anxiety disorders and attachment insecurity)
 - children and young people from particular black or minority ethnic groups
 - girls with a diagnosis of, or at risk of developing conduct disorders.

4.1.2 Groups that will not be covered

Recommendations will not be made specifically for the following groups, although parts of the guideline may be relevant to their care.

- a) Adults (aged 19 and older).
- b) Children and young people with psychosis.
- c) Children and young people with autism spectrum conditions.
- d) Primary drug and alcohol problems.
- e) Children and young people with learning disabilities (with an IQ of less than 70).
- f) Children and young people with speech and language difficulties.

4.2 *Health and social care setting*

- a) Primary, secondary and tertiary health and social care settings, the criminal justice system and forensic services, children services and educational settings.
- b) Other settings in which NHS and social care services are funded or provided, or where NHS or social care professionals are working in multi-agency teams.
- c) The guideline will also comment on and include recommendations about the interface with other sectors and such as education services, youth services, the criminal justice system and the voluntary sector.

4.3 *Areas to be considered*

4.3.1 Key issues that will be covered

- a) The behaviours, signs or symptoms that should prompt healthcare, education and social care professionals, and others working with

children and young people, to consider the presence of a conduct disorder.

- b) Validity, specificity and reliability of the components of diagnostic assessment after referral, including:
- the structure for assessment
 - diagnostic thresholds
 - assessment of risk.
- c) Psychosocial interventions, including:
- individual and group psychological interventions
 - parenting and family interventions (including family-based prevention models)
 - social care (including interventions for looked after children and young people), vocational, educational and community interventions, and work with peer groups
 - multisystemic interventions.
- d) Pharmacological interventions, including antipsychotics and medications for ADHD. Note that guideline recommendations will normally fall within licensed indications; exceptionally, and only if clearly supported by evidence, use outside a licensed indication may be recommended. The guideline will assume that prescribers will use a drug's summary of product characteristics to inform decisions made with individual patients.
- e) Physical interventions, such as diet.
- f) The organisation, coordination and delivery of care, and care pathways for the components of treatment and management. This will include transition planning and will be based on an ethos of multi-agency and multiprofessional working.

4.3.2 Issues that will not be covered

- a) Specific interventions for sexually abused or traumatised children and young people.
- b) Specific interventions for children and young people with significant speech and language difficulties.
- c) Coexisting conditions if conduct disorder is not a primary diagnosis.
- d) Preventive interventions for the general population.
- e) Setting based interventions, for example, school based interventions, for those who are not at significant risk of developing conduct disorders.

4.4 Main outcomes

- a) Antisocial behaviour at home and at school (including offending behaviour).
- b) Psychological, educational and social functioning as rated by the child or young person, professionals (including teachers) and parents or carers.

4.5 Economic aspects

The guideline will take into account both clinical and cost effectiveness when making recommendations involving a choice between alternative interventions. A review of the economic evidence will be conducted and analyses will be carried out as appropriate. The preferred unit of effectiveness will be the quality-adjusted life year (QALY), but a different unit of effectiveness may be used depending on the availability of appropriate clinical and utility data for children and young people with conduct disorders and associated antisocial behaviours. Costs considered will be from an NHS and personal social services (PSS) perspective in the main analyses, and a criminal justice perspective may also be considered. Further detail on the methods can be found in 'The guidelines manual' (see 'Further information').

4.6 Status

4.6.1 Scope

This is the consultation draft of the scope. The consultation dates are 12 January to 9 February 2011.

4.6.2 Timing

The development of the guideline recommendations will begin in April 2011.

5 Related NICE guidance

5.1 Published guidance

5.1.1 NICE guidance to be updated

This guideline will update and replace the following NICE guidance: [\[or\]](#)
Depending on the evidence, this guideline might update and replace parts of the following NICE guidance:

- [Title of appraisal]. NICE technology appraisal guidance [number] ([Year]). Available from [www.nice.org.uk/guidance/TA\[number\]](http://www.nice.org.uk/guidance/TA[number])

5.1.2 Other related NICE guidance

- Antisocial personality disorder. NICE clinical guideline 77 (2009). Available from www.nice.org.uk/guidance/CG77
- Attention deficit hyperactivity disorder (ADHD). NICE clinical guideline 72 (2008). Available from www.nice.org.uk/guidance/CG72
- Parent-training/education programmes in the management of children with conduct disorders. NICE technology appraisal guidance 102 (2006). Available from www.nice.org.uk/guidance/TA102

6 Further information

Information on the guideline development process is provided in:

- 'How NICE clinical guidelines are developed: an overview for stakeholders the public and the NHS'

- 'The guidelines manual'.

These are available from the NICE website

(www.nice.org.uk/GuidelinesManual). Information on the progress of the guideline will also be available from the NICE website (www.nice.org.uk).